

Increased growth and reduced summer drought limitation at the southern limit of *Fagus sylvatica* L., despite regionally warmer and drier conditions.

Hacket-Pain, Andrew J. ^{1,2} [corresponding author]

1. St. Catherine's College, Manor Road, Oxford, OX1 3UJ, UK

2. Fitzwilliam College, Storeys Way, Cambridge, CB3 0DG, UK

ajh220@cam.ac.uk

Friend, Andrew D. ³

3. Department of Geography, University of Cambridge, Downing Place, Cambridge, CB2 3EN, UK

adf10@cam.ac.uk

Abstract

Tree populations at the equatorward edge of their distribution are predicted to respond to increased temperature and drought with declining performance. Empirical studies of *Fagus sylvatica* L., one of the most studied tree species in Europe, have broadly supported these predictions. Using a network of tree ring chronologies from northern Greece, we showed that growth in populations of this species at their southeast distribution limit was limited by summer temperature and precipitation, particularly at low elevations. Furthermore, decadal periods of lower precipitation and higher temperature in the twentieth century were associated with multi-year growth depressions. However, since 1990, growth trends were positive across the network, despite continued dry and hot summer conditions. Growth trends were not correlated with either elevation or tree age. Additionally, correlations between growth and temperature and precipitation were weaker in recent decades. These results are consistent with another recent report from the Balkan Peninsula, and indicate that forests in this region may be more resistant to regional climate change than previously considered.

Keywords

Beech; Greece; Balkan; range edge; tree growth; growth trend

29 **Introduction**

30 Populations at the equator-ward or low-elevation edge of their species distribution are expected to
31 respond to increased temperatures and drought with a reduction in performance (Woodward, 1987). In
32 tree species, this response is expected to be evident in declining growth, increased mortality and reduced
33 recruitment (Jump *et al.*, 2010). Changes in demographic and subsequent distribution shifts have already
34 been observed at the trailing edge of many tree species distributions in response to recent climate change
35 (e.g. Beckage *et al.*, 2008). *Fagus sylvatica* L. (European beech) is a widely distributed species in Europe,
36 with great economic and biodiversity value (Gessler *et al.*, 2007), and has often been used as a model
37 species for investigating the current and future impact of climate change on forest ecosystems (e.g.
38 Kramer *et al.*, 2010). It is widely considered to have high sensitivity to growing season drought, based on
39 evidence from experimental and physiological studies (Robson *et al.*, 2012) , forest inventory data
40 (Seynave *et al.*, 2008), dendrochronological studies (Piovesan *et al.*, 2005) and statistical and process-
41 based models of distribution (Kramer *et al.*, 2010). Furthermore, drought stress is widely considered to
42 limit the distribution of *F. sylvatica* at its southern distribution limit, and subsequently the increase in
43 temperature and summer drought in southern Europe over the last half century (Giorgi, 2002) are
44 expected to have led to reduced performance in many of these southern populations (Fyllas and
45 Troumbis, 2009). In contrast, in northern Europe, beech is expected to have increased in competitiveness
46 and expanded into areas currently occupied by boreo-nemoral species as a consequence of increasing
47 survival and seed maturation success, and higher growth rates (Koca *et al.*, 2006). These trends are
48 expected to continue over the next century as climate continues to warm. It is important to test these
49 predictions as they are key to predicting future range shifts, and if species responses to climate change
50 differ from expectations this will have far-reaching consequences for forest carbon cycling, biodiversity
51 and forest-based economic activities.

52 Until recently, empirical studies have broadly supported predictions, with evidence of increased growth
53 and competitiveness reported in northern Europe (Bascietto *et al.*, 2004; Bolte *et al.*, 2010) and growth
54 declines and increased mortality reported at the southern distribution edge (Jump *et al.*, 2006; Piovesan

55 *et al.*, 2008). Recently however, the picture has become more complex, with climate warming in northern
56 Europe linked with evidence of negative impacts of climate change on forest growth, including declines
57 in growth (Aertsens *et al.*, 2014; Latte *et al.*, 2015). Additionally, a recent study has reported increased
58 tree growth in southern European beech forests, during the ongoing period of regional warming (Tegel
59 *et al.*, 2014). There is therefore a pressing need to improve understanding of the relationships between
60 tree growth and climate in this species, particularly at the southern distribution margin where beech
61 forests are understudied in comparison to central and northern Europe. Furthermore, we require
62 additional understanding of the response of growth in this region to recent changes in climate, which
63 have been characterised by increasing temperatures and summer drought stress (Giorgi, 2002). Tree
64 rings provide a valuable way to investigate such growth responses, allowing reliable assessment of
65 annual aboveground net primary production (Bascietto *et al.*, 2004). However, tree-rings contain
66 multiple biological and environmental signals necessitating careful analysis and interpretation (e.g.
67 Bowman *et al.*, 2013).

68 This study uses a new network of 21 ring chronologies from across northern Greece to investigate
69 evidence of the impacts of recent climate change on tree growth at the southeast distribution limit of this
70 species. We specifically address the following research questions:

- 71 1. What is the relationship between tree growth (ring width) and climate at the southeast
72 distribution limit of beech?
- 73 2. How have interannual and decadal variations in growth in beech responded to recent changes in
74 climate?
- 75 3. What do growth-climate relationships and growth trends within the network tell us about
76 ongoing and likely future response of this species to climate change?

77

78 **Material and methods**

79 **Study Area**

80 The study was conducted in northern Greece, in forests which represent the most south-easterly
81 populations of beech in Europe. Beech forms the treeline on many mountains (~1900 m a.s.l.), and at
82 lower elevations beech is typically replaced by mixed deciduous oak-dominated woodland (~800 m a.s.l.)
83 (Tsiripidis *et al.*, 2007). Research focused on an intensively sampled elevation transect on Mt. Vermio
84 (40.64°N, 21.95 °E), with eleven sampled sites extending from the high elevation treeline at ~1950 m a.s.l.
85 to the lowest elevation site at 880 m a.s.l. (Table 1 and Figure 1). Additionally, 10 further sites were
86 sampled across northern Greece to test the regional consistency of results (Table 1 and Figure 1). This
87 additional network of sites included a stand on Mt. Oxia (38.78°N, 21.97°E), the most extreme southeast
88 population of beech in Europe, and the most southerly occurrence in the Balkan peninsula (Tsiripidis *et*
89 *al.*, 2007).

90 Climate in this region is typically Mediterranean, with warm dry summers and cold wet winters, although
91 there is geographical and elevational variation (Figure S 1 and S 2). Changes in climate over the last
92 century are consistent with other regions in the Mediterranean, with warming during the first half of the
93 20th century, followed by a trend of cooling temperature from 1950-1970. Since the 1970s temperatures
94 have risen. Annual precipitation has no significant long-term trend, but summer precipitation was lower
95 during the 1940s and 1950s, and during the decade centred on 2000.

96 **Tree ring sampling**

97 At each site increment cores were extracted from 20-31 canopy dominant trees with two (occasionally
98 three) cores taken per tree from opposite directions and parallel to any slope. Sample preparation
99 followed standard dendrochronological methods, involving the mounting of cores, and then surface
100 sanding until individual ring boundaries were clearly visible. Cores were scanned at either 1600 or 2400
101 dpi (the latter when rings were particularly narrow), and ring widths for each core was measured using
102 the software Coorecorder v7.3 (Larsson, 2003). Initial cross-dating was conducted in CDendro v7.3
103 (Larsson, 2010), and then checked for each individual site using the standard dendrochronological

104 software COFECHA (Grissino-Mayer, 2001). Some cores could not be successfully measured or cross-
105 dated and these were excluded from subsequent analysis. Commonly used chronology statistics were
106 used to confirm the suitability of our chronologies for dendrochronological analysis (Grissino-Mayer,
107 2001) (Table 2), and PCA analysis was used to quantify the common signal between sites in the network.

108

109 **Tree ring processing**

110 Raw ring width chronologies typically contain low-frequency signals (i.e. decadal and longer), associated
111 with changes in tree age and size, canopy position and long-term changes in the abiotic environment. We
112 used two techniques in order to account for these effects. In the first, we removed low-frequency variance
113 from each individual tree chronology by detrending the ring width time-series using the R package `dplR`
114 (Bunn et al. 2012), individually fitting the raw ring width series from each core with a 32-year cubic spline
115 with a 50% frequency cut off. Dimensionless ring width indices were created for each core by dividing
116 the observed ring width by the spline, with individual indices then averaged to produce a mean ring width
117 index (RWI) for each sampled site, with a prewhitening procedure applied to account for autocorrelation
118 in the RWI time-series. The RWI captures high frequency (i.e. interannual) variation in tree growth. In
119 the second technique we converted raw ring width measurements into basal area increment (BAI) using
120 the formula:

$$121 \quad \text{BAI}_t = \pi(r_t^2 - r_{t-1}^2)$$

122 where r is the radius of the tree in year t . BAI chronologies account for the geometric effect (increasing
123 cambium area as tree size increases), but retain low frequency variation due to changes in growing
124 conditions (e.g. climate, canopy position). A mean site BAI chronology is expected to show a long-term
125 positive trend during the juvenile phase, partly due to stand dynamics and changes in tree size. The early
126 parts of a mean BAI chronology represent suppressed growth only, with increasing BAI in the mean site
127 chronology as sampled trees increase in size and gain access to the canopy (Bowman *et al.*, 2013). In
128 order to account for this potential bias, individual tree BAI chronologies were truncated to include only
129 mature-phase growth. The cut-off was determined as either the end of the most recent canopy release

130 event, or the transition to mature growth (following the method of Jump *et al.*, (2006), applied to
131 individual trees) (see Figure S 3 for more details). Individual chronologies were averaged for each site to
132 create mature BAI (mBAI) chronologies, and then scaled and centred for plotting and analysis. BAI
133 chronologies created without the removal of juvenile phase growth are included in the supplementary
134 information.

135 **Climate data**

136 Climate data were extracted from the gridded half-degree dataset CRU TS 3.10.01 (Harris *et al.*, 2013).
137 We used monthly-mean daily 24-hour maximum temperature (mean maximum, *MAX*) and monthly
138 precipitation (*PREC*). The location of each grid cell in relation to the sample sites is show in Figure 1. We
139 note that this climate dataset represents regional climates, and so will not fully capture the growing
140 condition experienced at the individual or population scale. Short datasets from high elevation
141 meteorological stations have recently been established by the National Observatory of Athens, and were
142 used for comparison with the gridded dataset (Figure S 1). In addition to temperature and precipitation
143 data, we also quantified drought using the Standardized Precipitation-Evapotranspiration Index (SPEI,
144 Vicente-Serrano *et al.*, 2010) using a 3-month window such that the drought index for July incorporated
145 data for May, June and July (with all months equally weighted). In addition to the gridded climate data,
146 regional climate reconstructions were used to provide information on regional climate trends prior to
147 the observational record (Griggs *et al.*, 2007; Klesse *et al.*, 2015; Levanic *et al.*, 2015).

148 **Ancillary data**

149 The diameter at breast height (DBH) and tree height were measured individually for all sampled trees
150 using a tape measure and a Haglof Vertex III hypsometer. For each sampled tree, age was calculated by
151 estimating the number of missing inner rings for each core (i.e. when the core missed the centre of the
152 tree) and adding this to the number of measured rings. The number of missing rings for each core was
153 estimated by dividing the distance from the innermost measured ring to the pith by the mean ring width
154 of the ten innermost measured rings.

155 **Statistical analysis**

156 To quantify the relationship between annual growth and climate, we calculated bootstrapped correlation
157 coefficients and response functions between the prewhitened RWI chronologies and monthly climate
158 data (*PREC* and *MAX*), using the `bootRes` package in R (Zang and Biondi, 2013). Correlations were
159 calculated for the period 1958-2010, which fully incorporates all site chronologies in the network (Table
160 2), and limits analysis to the period of more comprehensive climate station data in Greece (although note
161 that the gridded dataset CRU TS 3.2.10 was used for the analysis). Moving Correlation Analysis (MCA)
162 was used to assess the temporal stability of growth-climate relationships. 20 year intervals were used for
163 MCA, with a ten year time-step (i.e. a total of four windows). To assess changes in tree growth, and
164 compare changes in growth between sites, mBAI chronologies were standardised by subtracting the
165 chronology mean and dividing by the standard deviation, and linear trends for the period 1991-2010
166 calculated using linear regression.

167

168 **Results**

169 **Growth-climate relationships**

170 The growth-climate analysis of the 21 chronologies revealed relationships between *RWI* and both
171 temperature and precipitation for the period 1958-2010 (Figure 2 and S 4). The results of the correlation
172 and response function analyses were broadly consistent, and subsequent analysis therefore focused on
173 the response function analysis only, as it accounts for autocorrelation between monthly climate variables.
174 For temperature (*MAX*) the strongest signals negative relationships between *RWI* and temperature in the
175 summer of growth, and at some sites, with previous summer temperature (Figure 2). However, for the
176 year of growth the relationships were significant in the response function analysis in only four
177 chronologies (all July). In the correlation analysis, 13 chronologies had at least one significant correlation
178 between *RWI* and temperature for either June, July or August in the year of growth (Figure S4).
179 Relationships between *RWI* and summer temperature were stronger at low elevations, both along the
180 Vermio transect and in the rest of the network (Figure 2). At some high elevation sites growth was
181 favoured by warm temperatures in May (significant at two sites), but relationships between growth and
182 temperature were negative for the remaining months of the growing season at all sites.

183 Strong positive relationships between *RWI* and precipitation were found at sites across the network, with
184 precipitation in the growing season and previous summer the key factors (Figure 2). The strength of these
185 relationships also showed strong variation with elevation throughout the early- to mid-summer (May-
186 July), with strong and frequently significant positive relationships at low elevation sites. At higher
187 elevations, relationships were weaker, but only rarely negative (Figure 2). However, *RWI* was negatively
188 related to precipitation in April, especially at low elevation sites along the Vermio transect. *RWI* was
189 generally positively related to precipitation in the late summer prior to the growing season (July-
190 September), with significant response functions at six high or medium elevation sites.

191 **Growth-climate relationships through time**

192 Testing the stability of growth-climate relationships was challenging due to the relatively short length of
193 reliable climate records in this region, with CRU TS 3.10.01 based on increasingly sparse station data

194 prior to 1958. However, moving correlation analysis (MCA) showed that the strength of growth-climate
195 relationships varied through time during the period 1961-2010 (Figure 3). Overall, correlations between
196 RWI and summer temperature (MAX_{JJA}) were negative across all 21 sites, but generally became weaker
197 in recent decades (i.e. the 1981-2000 and 1991-2010 windows), particularly at low elevations along the
198 Vermio transect (although relationships were rarely significant). For example, along the Vermio transect,
199 5 of the 11 sites had significant negative correlations between RWI and MAX_{JJA} for the window 1971-1990,
200 but for the most recent window 1991-2010 only one of these correlations was still significant, and the
201 correlation coefficient had declined for 10 of the sites.

202 For precipitation, the most recent window was also associated with an apparent a decline in the strength
203 of correlations (Figure 3). This decline in correlation was stronger for sites on the Vermio transect, where
204 the correlation coefficients shifted to <0.2 (i.e. close to zero) for the 1991-2010 window for all but one
205 site. Correlations were >0.2 for at least eight sites for all the previous three windows. Similar shifts in
206 correlations were found in the rest of the network, especially at low and medium elevations.

207 **Long-term growth trends**

208 Throughout most of the twentieth century, decadal trends in mBAI broadly mirrored decadal trends in
209 summer drought stress with lower growth during periods with drier and warmer summers (Figure 4).
210 This was particularly evident along the Vermio transect. For example, the periods 1920-1930 and 1945-
211 1955 were associated with lower mBAI at most sites. However, recent mBAI trends were positive across
212 the network, including at high and low elevations, and at the most southerly sites (Figure 4). Importantly,
213 trends for the period 1991-2010 were not significantly correlated with the age of the sampled trees
214 (mean sampled tree age). Additionally, in all but two cases, trends for the period 1991-2010 were more
215 positive than the period 1971-1990, indicating that positive trends were not due to age effects (Table S
216 1). Furthermore, the decade centred on the year 2000, which had consistently low summer precipitation
217 was generally not associated with a strong growth suppressions, in contrast to previous periods with
218 similar climate conditions. Indeed, at most sites mBAI showed strongly positive trends through this
219 period (significant at seven of the eleven sites on Mt Vermio, and at two of the ten sites in the rest of the

220 network). Only three sites showed evidence of negative mBAI trends in the period 1991-2010; these were
221 two treeline site (VERM19 and VERMTL) and the lowest elevation site in the network (OSSA06), although
222 at both these sites the trends were close to zero and not statistically significant. Consequently, there was
223 no overall significant relationship between mBAI trend and elevation across the network, and with the
224 exception of OSSA06 all low elevation sites had positive trends in mBAI.

225

Discussion

Response of growth to recent changes in climate

The response function analysis revealed summer temperature and precipitation control on interannual variations in growth, with strong and frequently significant relationships between RWI and summer temperature and precipitation across the network (Figure 2). The relationships were particularly strong at low elevations. The association between mBAI and climate also indicated drought limitation on growth with mBAI decreases associated with dry (and hot) periods in the 1920s, during the 1940s and 1950s, and to a more limited extent during the 1980s (Figure 4). These results were consistent along the Vermio transect (11 sites) and, to a more limited extent, across the rest of northern Greece (10 sites). Previous studies have reported strong coherence between beech growth trends in southern Europe and decadal changes in climate (Piovesan *et al.*, 2008; Castagneri *et al.*, 2014; Tegel *et al.*, 2014). Together with the results of the response function analysis, these results have important implications for the future of this species in southern Europe. If climate continues to warm without an increase in precipitation (Giorgi and Lionello, 2008), this will be expected to lead to a general reduction in the growth of beech at all elevations in Greece (although the response at the population level will be determined by a combination of regional climate changes and local factors, such as soil depth or nutrients). We see evidence to support this in our network, with declines in mBAI during the late 1970s and 1980s in many sites. However, since 1990 the association between growth trends and climate has weakened. For example, mBAI increased at most sites from around ~1990, despite many parts of the study region experiencing decadal-scale drought through the late 1990s and early 2000s (Figure 4). Additionally, MCA revealed weaker correlations between RWI and summer temperature and precipitation in recent decades (and specifically for the period associated with these positive mBAI trends, 1991-2010). The positive mBAI trends, and the apparent decoupling of both decadal growth trends and RWI from regional summer climate, are consistent with some recent studies of beech elsewhere in southern Europe (Tegel *et al.*, 2014), but contradict other tree ring studies and also model simulations of growth declines in low elevation beech forests (Jump *et al.*, 2006; Piovesan *et al.*, 2008; Fyllas and Troumbis, 2009). Furthermore, while Tegel *et al.* (2014) also reported strong

recent increases in beech growth in the Balkan region, these forests were located in regions of higher summer precipitation, while the sites in this network are located in the drier southeast of the Balkan peninsula, close to the xeric limit of the species distribution. Additionally, while Cavin and Jump (2016) found little evidence of growth declines in southern beech forests in the Iberian peninsula, these sites were also associated with weak growth-climate correlations. This can be interpreted as indicating that these sampled trees represent “climate-relics”, occupying specific locations where microclimates dominate (Cavin and Jump, 2016). In contrast, the sites in the Greek network reported in this study show stronger relationships between *RWI* and regional summer temperature and precipitation, indicating coupling of forest growth to regional weather conditions. Consequently, these forests are expected to also respond to decadal changes in regional climate, including the recent increases in summer temperature.

The lack of evidence for a decline in recent growth in Greek beech populations over recent decades, despite increasing drought stress could result from biases in tree-ring chronologies (Bowman *et al.*, 2013), but these are unlikely to fully explain the positive trends we detect, particularly as we account for a major source of bias by considering only mature-phase growth (*mBAI*). Alternatively, the recent increases in *BAI* across the whole elevation gradient could be the result of environmental changes such as increasing atmospheric CO_2 , reduction in air pollutants including ozone, or changes in management. Carbon assimilation in Greek beech forests is strongly limited by stomatal conductance during dry conditions (Raftoyannis and Radoglou, 2002), and an increase in water use efficiency (*WUE*) as a consequence of higher atmospheric CO_2 will be expected to lead to increased growth in a carbon-limited ecosystem (assuming negligible effects of any changes in humidity). However, numerous studies have failed to detect growth increases, despite increased *WUE*, including in other southern European beech forests (e.g. Penuelas *et al.*, 2008).

Relationships between growth and climate

The analysis of relationships between growth and regional climate revealed that summer temperature and precipitation were the key climate signals in beech tree ring chronologies from across Greece (Figure 4). Summer drought has been identified as the key driver of beech tree-ring chronologies across the

species distribution of beech, including at the southern distribution edge (Dittmar *et al.*, 2003; Piovesan *et al.*, 2005; Hacket Pain *et al.*, 2016). In the only other published investigations of beech chronologies from the Balkan peninsula, Tegel *et al.* (2014) and Castagneri *et al.* (2014) also found that growth of beech was sensitive to summer climate, but that temperature was a stronger control than precipitation. However both of these studies noted the poor quality of climate data in the western Balkans, which likely influenced the strength of statistical relationships. Additionally, annual and summer precipitation is much higher in mountains of the western Balkans than in the region studied here (Figure S 2). While growth was limited by summer temperature and precipitation across the network in this study, the strength of the relationships were stronger at the lowest elevation sites (Figure 2). This variation in response function coefficients with elevation was particularly strong for summer precipitation, consistent with numerous previous studies of beech (e.g. Hartl-Meier *et al.*, 2014), and indicates that growth at low elevation sites is more strongly controlled by water availability. The low elevation sites in this study represent trees growing close to the distribution limit, which according to previous studies and the classic dendrochronological literature are expected to show stronger relationships between growth and summer drought (Fritts, 1966).

However, while relationships between growth and summer climate were strongest at low elevations, the strength of the negative relationships with temperature at high elevations, including at the treeline, was surprising (although note that these were only ever significant in teh correlation analysis, Fig S 4). Low temperatures are commonly thought to limit tree growth in high elevation or latitude forests, and tree-ring studies have tended to find a positive effect of higher temperatures on growth at high elevations (Wettstein *et al.*, 2011), including in beech forests (Dittmar *et al.*, 2003; Latte *et al.*, 2015). However, Piovesan *et al.*, (2005) found strong summer drought signals in high elevation beech forests in Italy, and recent studies from conifer treelines in the Balkans have found similar results (Panayotov *et al.*, 2010; Klesse *et al.*, 2015). This has been linked to thin soils and steep slopes at high elevations, which also applies in this study (Supplementary Table 1).

304 Strong and significant relationships between growth and the temperature and precipitation of the
305 previous summer were also common across the network (Figure 4 and S4), but were strongest at highest
306 elevations and in the sites with the oldest trees. Unfortunately, as the site elevation and mean site tree
307 age are correlated in our network, it is difficult to distinguish these two influences, although future
308 analysis at the individual tree level may allow separation of age and elevation effects. In conifers, the
309 importance of previous summer weather in treeline chronologies has been linked to the short growing
310 season, which reduces the availability of carbohydrates at the start of the following growing season
311 (Panayotov *et al.*, 2010). Similarly, in beech, 20-30% of annual growth is thought to rely on carbon
312 assimilated in the previous year (Cufar *et al.*, 2008). An alternative explanation for the correlations
313 between growth and previous summer climate is related to masting phenomenon (Hacket-Pain *et al.*,
314 2015). Years of heavy seed production are cued by conditions in the previous summer, and are
315 themselves associated with strong growth reductions (Drobyshev *et al.*, 2010). The increase in the
316 strength of correlations with previous summer temperature with increased elevation (and age) might
317 result from greater growth reductions in mast years, due to stronger growth-reproduction trade-offs at
318 sites with longer growing seasons or with older trees (Thomas, 2011). These hypotheses are difficult to
319 test in this study due to the lack of data on seed production in the Balkan region, but two mast years
320 reported in beech forests in northern Greece by Gerasimidis *et al.* (2006) correspond to years of low
321 growth across the network (1998 and 2001).

322 **Conclusions**

323 We found inter-annual growth of *F. sylvatica*, as estimated using tree-ring chronologies, in southeastern
324 Europe to be correlated with summer precipitation and temperature. Trees at all elevations were
325 sensitive to summer drought, including sites at the treeline. Until the end of the 1990s, decadal trends in
326 mBAI closely matched regional summer climate, with growth decreases during periods of drier and
327 warmer summers. However, in recent decades relationships between inter-annual and decadal climate
328 variability and growth have weakened. Recent mBAI trends are positive across the network, including at
329 low elevation sites, despite an ongoing warming trend in the region. These results were consistent along
330 both an intensively sampled elevation transect (Mt. Vermio) and in a network of sites across northern
331 Greece. We suggest that these results indicate that beech forests in this region may be more resistant to
332 regional climate changes than previously considered.

333




Figure 1. The location of the sites included in the *F. sylvatica* tree ring network in Greece. The rectangle labelled VERMIO shows the location of the transect on Mt. Vermio, enlarged in the inset map. The dashed boxes represent the grid cells for the gridded climate data (CRU TS 3.10.01). Note that the two lowest elevation sites in the Vermio transect (VERM10 and VERM08) fall slightly outside grid cell, but the decision was taken to use the same climate data as the rest of the transect for consistency.




Figure 2. Response function coefficients of ring width index (*RWI*) and monthly climate variables for each site. a) Monthly mean maximum temperature (*MAX*). b) Monthly total precipitation (*PRE*). Response functions were calculated for the period 1958-2010 (common to all sites) and significance was calculated using bootstrapping (95% confidence level), and indicated by black circles. Sites are ordered by elevation.





Figure 3. Moving Correlation Analysis, calculated using 20 year windows with 10 year time-steps, and plotted according to the end year of each window. Filled points indicate significance at the 95% confidence level, estimated using bootstrapping. a) MAX_{JJA} for the Vermio transect. b) PRE_{MJJ} for the Vermio transect. c) MAX_{JJA} for the rest of the network. d) PRE_{MJJ} for the rest of the network.



352

353 **Figure 4.** Mature phase standardised Basal Area Increment (mBAI) chronologies for each site. a) mBAI
 354 chronologies for sites on the Vermio transect, ordered vertically by elevation (with an cumulative offset
 355 of 2 SD units). Splines of mBAI chronologies are used to highlight growth trends (frequency response =
 356 50%, wavelength = 16 years). Sites are ordered by elevation. The inset plot shows the slope of the linear
 357 trend in mBAI over the period 1991-2010, plotted against the elevation of the site. Significant slopes
 358 ($p < 0.05$) are plotted in black. b) as for a), but for the rest of the network. c) CRU TS 3.10.01 climate data
 359 for the period 1901-2010, with splines plotted to highlight trends. Climate data was scaled and centred.

Acknowledgements

We thank Dimitris Zianis for help organising access to field sites in Greece, and to local forestry authorities for permission to conduct fieldwork. We are grateful to T. Rademacher, N. Peters, N. Evans, J. Rae and M. Tucker for assisting with field data collection. The National Observatory of Athens provided climate data.

Funding

AHP received funding from the Department of Geography, University of Cambridge and Clare College, Cambridge.

References

- Aertsen, W., Janssen, E., Kint, V., Bontemps, J.D., Van Orshoven, J., Muys, B., 2014. Long-term growth changes of common beech (*Fagus sylvatica* L.) are less pronounced on highly productive sites. *Forest Ecology and Management* 312, 252-259.
- Bascietto, M., Cherubini, P., Scarascia-Mugnozza, G., 2004. Tree rings from a European beech forest chronosequence are useful for detecting growth trends and carbon sequestration. *Canadian Journal of Forest Research-Revue Canadienne De Recherche Forestiere* 34, 481-492.
- Beckage, B., Osborne, B., Gavin, D.G., Pucko, C., Siccama, T., Perkins, T., 2008. A rapid upward shift of a forest ecotone during 40 years of warming in the Green Mountains of Vermont. *Proceedings of the National Academy of Sciences of the United States of America* 105, 4197-4202.
- Bolte, A., Hilbrig, L., Grundmann, B., Kampf, F., Brunet, J., Roloff, A., 2010. Climate change impacts on stand structure and competitive interactions in a southern Swedish spruce-beech forest. *European Journal of Forest Research* 129, 261-276.
- Bowman, D.M.J.S., Brien, R.J.W., Gloor, E., Phillips, O.L., Prior, L.D., 2013. Detecting trends in tree growth: not so simple. *Trends in Plant Science* 18, 11-17.
- Castagneri, D., Nola, P., Motta, R., Carrer, M., 2014. Summer climate variability over the last 250 years differently affected tree species radial growth in a mesic *Fagus-Abies-Picea* old-growth forest. *Forest Ecology and Management* 320, 21-29.
- Cavin, L., Jump, A.S., 2016. Highest drought sensitivity and lowest resistance to growth suppression are found in the range core of the tree *Fagus sylvatica* L. not the equatorial range edge. *Global Change Biology*.
- Cufar, K., Prisan, P., de Luis, M., Gricar, J., 2008. Tree-ring variation, wood formation and phenology of beech (*Fagus sylvatica*) from a representative site in Slovenia, SE Central Europe. *Trees-Structure and Function* 22, 749-758.
- Dittmar, C., Fricke, W., Elling, W., 2006. Impact of late frost events on radial growth of common beech (*Fagus sylvatica* L.) in Southern Germany. *European Journal of Forest Research* 125, 249-259.
- Dittmar, C., Zech, W., Elling, W., 2003. Growth variations of Common beech (*Fagus sylvatica* L.) under different climatic and environmental conditions in Europe - a dendroecological study. *Forest Ecology and Management* 173, 63-78.
- Drobyshev, I., Overgaard, R., Saygin, I., Niklasson, M., Hickler, T., Karlsson, M., Sykes, M.T., 2010. Masting behaviour and dendrochronology of European beech (*Fagus sylvatica* L.) in southern Sweden. *Forest Ecology and Management* 259, 2160-2171.
- Fritts, H.C., 1966. Growth-rings of trees: their correlation with climate. *Science* 154, 973-979.
- Fyllas, N.M., Troumbis, A.Y., 2009. Simulating vegetation shifts in north-eastern Mediterranean mountain forests under climatic change scenarios. *Global Ecology and Biogeography* 18, 64-77.
- Gerasimidis, A., Panajiotidis, S., Hicks, S., Athanasiadis, N., 2006. An eight-year record of pollen deposition in the Pieria mountains (N. Greece) and its significance for interpreting fossil pollen assemblages. *Review of Palaeobotany and Palynology* 141, 231-243.
- Gessler, A., Keitel, C., Kreuzwieser, J., Matyssek, R., Seiler, W., Rennenberg, H., 2007. Potential risks for European beech (*Fagus sylvatica* L.) in a changing climate. *Trees-Structure and Function* 21, 1-11.
- Giorgi, F., 2002. Variability and trends of sub-continental scale surface climate in the twentieth century. Part I: observations. *Climate Dynamics* 18, 675-691.
- Giorgi, F., Lionello, P., 2008. Climate change projections for the Mediterranean region. *Global and Planetary Change* 63, 90-104.
- Griggs, C., DeGaetano, A., Kuniholm, P., Newton, M., 2007. A regional high-frequency reconstruction of May-June precipitation in the north Aegean from oak tree rings, AD 1089-1989. *International Journal of Climatology* 27, 1075-1089.
- Grissino-Mayer, H.D., 2001. Crossdating accuracy: A manual and tutorial for the computer program COFECHA. *Tree-Ring Research* 57, 205-221.

423 Hackett-Pain, A.J., Cavin, L., Friend, A.D., Jump, A., 2016. Consistent limitation of growth by high temperature
 424 and low precipitation from range core to southern edge of European Beech indicates widespread vulnerability
 425 to changing climate. *European Journal of Forest Science*.

426 Hackett-Pain, A.J., Friend, A.D., Lagueard, J.G.A., Thomas, P.A., 2015. The influence of masting phenomenon on
 427 growth-climate relationships in trees: explaining the influence of previous summers' climate on ring width.
 428 *Tree Physiology* 35, 319-330.

429 Harris, I., Jones, P.D., Osborn, T.J., Lister, D.H., 2013. Updated high-resolution grids of monthly climatic
 430 observations - the CRU TS3.10 Dataset. *International Journal of Climatology*.

431 Hartl-Meier, C., Dittmar, C., Zang, C., Rothe, A., 2014. Mountain forest growth response to climate change in
 432 the Northern Limestone Alps. *Trees-Structure and Function* 28, 819-829.

433 Jump, A.S., Cavin, L., Hunter, P.D., 2010. Monitoring and managing responses to climate change at the
 434 retreating range edge of forest trees. *Journal of Environmental Monitoring* 12, 1791-1798.

435 Jump, A.S., Hunt, J.M., Penuelas, J., 2006. Rapid climate change-related growth decline at the southern range
 436 edge of *Fagus sylvatica*. *Global Change Biology* 12, 2163-2174.

437 Klesse, S., Ziehmer, M., Rousakis, G., Trouet, V., Frank, D., 2015. Synoptic drivers of 400 years of summer
 438 temperature and precipitation variability on Mt. Olympus, Greece. *Climate Dynamics* 45, 807-824.

439 Koca, D., Smith, B., Sykes, M.T., 2006. Modelling regional climate change effects on potential natural
 440 ecosystems in Sweden. *Climatic Change* 78, 381-406.

441 Kramer, K., Degen, B., Buschbom, J., Hickler, T., Thuiller, W., Sykes, M.T., de Winter, W., 2010. Modelling
 442 exploration of the future of European beech (*Fagus sylvatica* L.) under climate change-Range, abundance,
 443 genetic diversity and adaptive response. *Forest Ecology and Management* 259, 2213-2222.

444 Larsson, L.-A., 2003. Coo-Recorder: image co-ordinate recording program. In, Available at <http://cybis.se>.

445 Larsson, L.-A., 2010. CDendro. In. Cybis Elektronik and Data AB, Saltsjöbaden, Sweden.

446 Latte, N., Lebourgeois, F., Claessens, H., 2015. Increased tree-growth synchronization of beech (*Fagus sylvatica*
 447 L.) in response to climate change in northwestern Europe. *Dendrochronologia* 33, 69-77.

448 Levanic, T., Poljansek, S., Toromani, E., 2015. Early summer temperatures reconstructed from black pine (*Pinus*
 449 *nigra* Arnold) tree-ring widths from Albania. *Holocene* 25, 469-481.

450 Panayotov, M., Bebi, P., Trouet, V., Yurukov, S., 2010. Climate signal in tree-ring chronologies of *Pinus peuce*
 451 *and Pinus heldreichii* from the Pirin Mountains in Bulgaria. *Trees-Structure and Function* 24, 479-490.

452 Penuelas, J., Hunt, J.M., Ogaya, R., Jump, A.S., 2008. Twentieth century changes of tree-ring delta C-13 at the
 453 southern range-edge of *Fagus sylvatica*: increasing water-use efficiency does not avoid the growth decline
 454 induced by warming at low altitudes. *Global Change Biology* 14, 1076-1088.

455 Piovesan, G., Biondi, F., Bernabei, M., Di Filippo, A., Schirone, B., 2005. Spatial and altitudinal bioclimatic zones
 456 of the Italian peninsula identified from a beech (*Fagus sylvatica* L.) tree-ring network. *Acta Oecologica-*
 457 *International Journal of Ecology* 27, 197-210.

458 Piovesan, G., Biondi, F., Di Filippo, A., Alessandrini, A., Maugeri, M., 2008. Drought-driven growth reduction in
 459 old beech (*Fagus sylvatica* L.) forests of the central Apennines, Italy. *Global Change Biology* 14, 1265-1281.

460 Raftoyannis, Y., Radoglou, K., 2002. Physiological responses of beech and sessile oak in a natural mixed stand
 461 during a dry summer. *Annals of Botany* 89, 723-730.

462 Robson, T.M., Sanchez-Gomez, D., Cano, F.J., Aranda, I., 2012. Variation in functional leaf traits among beech
 463 provenances during a Spanish summer reflects the differences in their origin. *Tree Genetics & Genomes* 8,
 464 1111-1121.

465 Seynave, I., Gegout, J.C., Herve, J.C., Dhote, J.F., 2008. Is the spatial distribution of European beech (*Fagus*
 466 *sylvatica* L.) limited by its potential height growth? *Journal of Biogeography* 35, 1851-1862.

467 Tegel, W., Seim, A., Hakelberg, D., Hoffmann, S., Panev, M., Westphal, T., Buntgen, U., 2014. A recent growth
 468 increase of European beech (*Fagus sylvatica* L.) at its Mediterranean distribution limit contradicts drought
 469 stress. *European Journal of Forest Research* 133, 61-71.

470 Thomas, S.C., 2011. Age-Related Changes in Tree Growth and Functional Biology: The Role of Reproduction.
 471 *Size- and Age-Related Changes in Tree Structure and Function* 4, 33-64.

472 Tsiripidis, I., Bergmeier, E., Dimopoulos, P., 2007. Geographical and ecological differentiation in Greek *Fagus*
 473 forest vegetation. *Journal of Vegetation Science* 18, 743-750.

474 Vicente-Serrano, S.M., Begueria, S., Lopez-Moreno, J.I., 2010. A Multiscalar Drought Index Sensitive to Global
475 Warming: The Standardized Precipitation Evapotranspiration Index. *Journal of Climate* 23, 1696-1718.
476 Wettstein, J.J., Littell, J.S., Wallace, J.M., Gedalof, Z.e., 2011. Coherent Region-, Species-, and Frequency-
477 Dependent Local Climate Signals in Northern Hemisphere Tree-Ring Widths. *Journal of Climate* 24, 5998-6012.
478 Woodward, F.I., 1987. *Climate and Plant Distribution*. Cambridge University Press, Cambridge.
479 Zang, C., Biondi, F., 2013. Dehydroclimatic calibration in R: The bootRes package for response and correlation
480 function analysis. *Dendrochronologia* 31, 68-74.

481